

On non- acceptability of variational based calculation in PT symmetrized complex Harmonic Oscillators : Direct study.

Biswanath Rath

Department of Physics, North Orissa University, Takatpur, Baripada -757003,
Odisha, INDIA

We notice through a direct calculation that any variational based calculation on PT symmetrized complex Harmonic oscillator can lead to break down of PT symmetry condition on real spectra .Two different types of oscillators have been tested yielding an uniform conclusion.

PACS 03.65Db,11.30.Er

Key words- PT symmetry, complex oscillator ,variational method ,broken PT symmetry.

I.Introduction

Study of Harmonic Oscillator characterized by the Hamiltonian

$$H = p^2 + W^2 x^2 \quad (1)$$

is interesting not only because of its exact solvability nature[1] displaying the energy eigenvalue

$$E_n = (2n + 1)W \quad (2)$$

and wave function

$$\Psi_n(x) = \left[\frac{\sqrt{W}}{\sqrt{\pi n! 2^n}} \right]^{1/2} H_n(\sqrt{W}x) e^{-\frac{Wx^2}{2}} \quad (3)$$

but also a testing laboratory for any new approximation[2,3] when it is under the influence of suitable non-Hermitian transformation in its momentum or co-ordinate .It is commonly believed that as long as the transformation is linear in nature i.e

$$p \rightarrow p' = p + iLx = y \quad (4a)$$

or

$$x \rightarrow x' = x + iRp = z \quad (4b)$$

the energy eigenvalue of the system remains invariant ,which has been proved in a similar problem by Ahmed [4] using gaugelike transformation , Rath and Mallick[5] using simple perturbation theory and Fernandez [6] using rigorous similarity transformation .Further Bender ,Hook and Klevansky[7] noticed negative energy eigenvalue in complex plane .Fernandez [6] also proved the similar thing using similarity transformation .However Rath [8] noticed the negative energy [8] in a similar problem under the infuence of real wave function . Further Rath[8] has noticed for certain typical constants , non- Hermitian Harmonic Oscillator can exhibit coplex energy .However author does not give any systematic study of complex nature of eigenvalue. It is important to note that once the transformation is taken in to consideration ,the transformed Hamiltonian

$$H = (p + iLx)^2 + W^2x^2 \quad (5)$$

changes its behaviour from hermiticity[1] to non-Hermiticity [1-3]. Hence the standard hermiticity [1] property i.e

$$H = H^+ \quad (6)$$

fails to say any thing about its energy eigenvalue and wave function. However ,the commutation relation

$$[x, p + iLx] = [x, y] = i \quad (7a)$$

or

$$[x + iRp, p] = [z, p] = i \quad (7b)$$

invariance demands the preservation of its energy eigenvalue i.e iso-spectral characteristics with the original Hamiltonian . The question now comes to reader's mind regarding the nature of energy eigenvalue .To support this we take the help of PT symmetry idea of Bender and Boettecher [2] or its mofied version the pseudo-Hermiticity [3] condition i.e

$$\eta H \eta^{-1} = H^+ \quad (8)$$

on real spectra ,which has been used previously by Ahmed [4] in a diffent form of Oscillator .A close observation on Eq(5) reflects that there are two parameters involved i.e L and W .Further any oscillator Hamiltonian like the one considered here (Eq(5)) has its own frequency of vibration (w) . Now question arrises how the frequency of vibration w is related with W, L in reflecting iso-spectral behaviour or negative spectrum and more interestingly when deviation from these two occurs .In other words for which values of W, L the system does not reflect real energy eigenvalue . Hence the aim of this paper is to make a direct study on the validity or failure of PT symmetric condition on real spectra by visualising it in Harmonic Oscillators under the influence of co-ordinate or momentum transformation with real wave function having positive frequency of vibration $w > 0$.

II. Non-Hermitian Oscillator in second quantization notation.

The Hamiltonian

$$H = (p + iLx)^2 + W^2 x^2$$

in second quantization can be expressed by replacing the co-ordinate and momentum as :

$$x = \frac{(a + a^+)}{\sqrt{2w}} \quad (9a)$$

and

$$p = i\sqrt{\frac{w}{2}}(a^+ - a) \quad (9b)$$

where the creation operator, a^+ and annihilation operator a satisfy the commutation relation

$$[a, a^+] = 1 \quad (10)$$

and w is an unknown parameter. Further in number space we have

$$a|n\rangle_w = \sqrt{n}|n-1\rangle_w \quad (10a)$$

$$a^+|n\rangle_w = \sqrt{(n+1)}|n+1\rangle_w \quad (10b)$$

$$\langle n|a^+a|n\rangle_w = n \quad (10c)$$

where the state vector $|n\rangle_w$ can be expressed as

$$|n\rangle_w = \left[\frac{\sqrt{w}}{\sqrt{\pi n!} 2^n}\right]^{1/2} H_n(\sqrt{w}x) e^{-\frac{wx^2}{2}} \quad (11)$$

Here we express H as

$$H = H_D + H_N \quad (12a)$$

where

$$H_D = \left[w + \frac{(W^2 - L^2)}{w}\right] \frac{(2a^+a + 1)}{2} \quad (12b)$$

and

$$H_N = U \frac{a^2}{2} + V \frac{(a^+)^2}{2} \quad (13c)$$

with

$$V = \left[-w + \frac{(W^2 - L^2)}{w} - 2L\right] \quad (13d)$$

$$U = \left[-w + \frac{(W^2 - L^2)}{w} + 2L\right] \quad (13e)$$

III. Parameter selection using perurbation theory .

(IIIA) Parameters added i.e $w_1 = L + W$.

Here we adopt effective way of determining Hamiltonian as suggested by Nesbet [] by making the coefficient of a^2 to zero i.e

$$U = [-w + \frac{(W^2 - L^2)}{w} + 2L] = 0 \quad (14a)$$

On solving the relation we get ,

$$w = w_1 = L + W \quad (14b)$$

In this case using the Eq(11) with $w = w_1 > 0$ in Eq(13b) we get

$$\langle n | H_D | n \rangle_{w_1} = E_n = (2n+1)W \quad (14c)$$

(IIIB) Parameters subtracted i.e $w_2 = L - W > 0$.

Similarly from $U = 0$ we get the frequency

$$w = w_2 = L - W > 0 \quad (14d)$$

and we get

$$\langle n | H_D | n \rangle_{w_2} = E_n = -(2n+1)W \quad (14e)$$

(IIIC) Parameters subtracted i.e $w_3 = W - L > 0$. Now from $V = 0$ we get the frequency

$$w = w_3 = W - L > 0 \quad (14f)$$

and we get

$$\langle n | H_D | n \rangle_{w_3} = E_n = (2n+1)W \quad (14g)$$

IV. Parameter selection using variational method .

In the above we have determined the parameter w from the non-diagonal terms however here we determine the parameter from the diagonal term using variational method as follows .

$$\epsilon_n = \langle n | H_D | n \rangle_w = [w + \frac{(W^2 - L^2)}{w}] \frac{(2n+1)}{2} \quad (15a)$$

Using variational method [] i.e

$$\frac{d\epsilon}{dw} = 0 \quad (15b)$$

we get

$$w = w_v = \sqrt{(W^2 - L^2)} \quad (15c)$$

V. Direct calculation of energy using Matrix Diagonalisation Method [MATLAB].

The MATLAB programme is given below as

B – MATLAB

$$N = 100; \quad (16a)$$

$$s = 1; \quad (16b)$$

$$n = 1 : N - 1; \quad (16c)$$

$$m = \sqrt{(n)}; \quad (16d)$$

$$W = 10; \quad (16e)$$

$$L = 5.4; \quad (16f)$$

$$w = \sqrt{(W^2 - L^2)}; \quad (16g)$$

$$x = \frac{s}{\sqrt{(2w)}}(diag(m, -1) + diag(m, 1)); \quad (16h)$$

$$p = \frac{is}{\sqrt{(2/w)}}(diag(m, -1) - diag(m, 1); \quad (16i)$$

$$y = (p + iLx); \quad (16j)$$

$$H = y^2 + W^2x^2; \quad (16k)$$

$$EigSort = sort(eig(H)); \quad (16l)$$

$$EigSort(1 : 40) \quad (16m)$$

VI. Complex Oscillator having Co-ordinate transformation

In this case we consider a different type of oscillator having non-Hermitian transformation in its co-ordinate as

$$H = L^2 p^2 + (x + iRp)^2 \quad (17)$$

Following the above procedure, here we just quote the key calculation of the following. The energy eigenvalue with $Q = 0$ can be written as

$$E_n = (2n + 1)L \quad (18)$$

Using perturbation theory we get only three values of w i.e

$$w_1 = \frac{1}{L - R} \quad (18a)$$

$$w_2 = \frac{1}{R + L} \quad (18b)$$

and

$$w_3 = \frac{1}{R - L} \quad (18c)$$

As per perturbative calculation we get iso-spectral nature

$$E_n(w_{1,2}) = (2n + 1)L \quad (19a)$$

and negative real spectrum

$$E_n(w_3) = -(2n + 1)L \quad (19b)$$

Now using variational method we get

$$w_4 = \frac{1}{\sqrt{(L^2 - R^2)}} \quad (20)$$

VII. Results and Conclusion.

In Table-I we present energy eigenvalues of complex PT symmetric oscillator for different values of L, W and w . It is observed that when the eigenvalues calculated

using frequency of oscillation $w = w_v$, we observe that system reflects real as well as complex eigenvalues. However when for the same values of L and W , if the frequency of oscillation is so adjusted that $w = W \pm L$ or $w = L - W$ we get real eigenvalues. It is interesting to note that for complex eigenvalues, the PT symmetry condition on real spectra of non-Hermitian Hamiltonian fails so also the pseudo-Hermiticity condition. For the second case present the result in Table-2. Here also we notice complex energy levels as in the first case and notice the broken PT symmetry in coordinate transformation. Lastly present systematic direct study reveals that under non-hermitian transformation system can reflect real (positive or negative) or complex energy depending on the choice of frequency of oscillation $w > 0$ thereby reflecting the validity or failure of PT symmetry condition.

References

- [1] L.I. Schiff, Quantum Mechanics, **3rd ed**(McGraw-Hill,Singapore, 1985);
- [2] C.M. Bender and S.Boettcher Phys.Rev.Lett.**80** ,5243 (1998).
- [3] A. Mostafazadeh arXiv :011016 [math-ph] .
- [4] Z. Ahmed, Phys. Lett. A **294**, 287 (2002).
- [5] B. Rath and P.Mallick arXiv ;1502.07891[quant-ph] (2015).
- [6] F.M.Fernandez,arXiv:1502.02694v2[quant-ph](2015)and
arXiv:submit/1200745[quant-ph](2015) .
- [7] C.M. Bender, D.W.Hook and S.P.Klevansky J.Phys **A 45** , 444003 (2012).
- [8] B. Rath, arXiv ;1502.07891[quant-ph],1505.05018[quant-ph] (2015).

Table -I

Energy eigenvalues of complex PT symmetric oscillator under momentum transformation with remarks .

W	L	w	Eigenvalues	$E_n = (2n + 1)W$ and Remarks
10	5.4	$\sqrt{W^2 - L^2}$	-10	few negative but real
10	5.4	$\sqrt{W^2 - L^2}$	-30	i.e $\rightarrow -E_n$
10	5.4	$\sqrt{W^2 - L^2}$	-50	
10	5.4	$\sqrt{W^2 - L^2}$	816.4 -28.4i	rest complex
10	5.4	$\sqrt{W^2 - L^2}$	816.4 +28.4i	PT condition fails
10	5.4	$W \pm L$	10	all positive
10	5.4	$W \pm L$	30	isospectral
10	5.4	$W \pm L$	50	with $E_n = (2n + 1)W$
10	5.4	$W \pm L$	70	
10	5.4	$W \pm L$	90	PT condition holds good
5.4	10	$L - W$	-5.4	all negative
5.4	10	$L - W$	-16.2	i.e $E_n = -(2n + 1)W$
5.4	10	$L - W$	-27.0	
5.4	10	$L - W$	-37.8	PT condition holds good

Table -II

Energy eigenvalues of complex PT symmetric oscillator under co-ordinate transformation with remarks .

R	L	w	Eigenvalues	$E_n = (2n + 1)L$ and Remarks
8	10	$\frac{1}{\sqrt{L^2 - R^2}}$	10	few positive but real
8	10		30	i.e $\rightarrow E_n$
8	10		50	
8	10		409.65-13.47i	rest complex
8	10		409.65 + 13.47i	PT condition fails
8	10	$\frac{1}{L \pm R}$	10	all positive
8	10		30	isospectral
8	10		50	with $E_n = (2n + 1)L$
8	10		70	
8	10		90	PT condition holds good
10	8	$\frac{1}{R-L}$	-8	all negative
10	8		-24	i.e $E_n = -(2n + 1)L$
10	8		-40	
10	8		-56	PT condition holds good

Table -I

Energy eigenvalues of complex PT symmetric oscillator under momentum transformation with remarks .

W	L	w	Eigenvalues	$E_n = (2n + 1)W$ and Remarks
10	5.4	$\sqrt{W^2 - L^2}$	-10	few negative but real
10	5.4	$\sqrt{W^2 - L^2}$	-30	i.e $\rightarrow -E_n$
10	5.4	$\sqrt{W^2 - L^2}$	-50	
10	5.4	$\sqrt{W^2 - L^2}$	816.4 -28.4i	rest complex
10	5.4	$\sqrt{W^2 - L^2}$	816.4 +28.4i	PT condition fails
10	5.4	$W \pm L$	10	all positive
10	5.4	$W \pm L$	30	isospectral
10	5.4	$W \pm L$	50	with $E_n = (2n + 1)W$
10	5.4	$W \pm L$	70	
10	5.4	$W \pm L$	90	PT condition holds good
5.4	10	$L - W$	-5.4	all negative
5.4	10	$L - W$	-16.2	i.e $E_n = -(2n + 1)W$
5.4	10	$L - W$	-27.0	
5.4	10	$L - W$	-37.8	PT condition holds good

Table -II

Energy eigenvalues of complex PT symmetric oscillator under co-ordinate transformation with remarks .

R	L	w	Eigenvalues	$E_n = (2n + 1)L$ and Remarks
8	10	$\frac{1}{\sqrt{L^2 - R^2}}$	10	few positive but real
8	10		30	i.e $\rightarrow E_n$
8	10		50	
8	10		409.65-13.47i	rest complex
8	10		409.65 + 13.47i	PT condition fails
8	10	$\frac{1}{L \pm R}$	10	all positive
8	10		30	isospectral
8	10		50	with $E_n = (2n + 1)L$
8	10		70	
8	10		90	PT condition holds good
10	8	$\frac{1}{R-L}$	-8	all negative
10	8		-24	i.e $E_n = -(2n + 1)L$
10	8		-40	
10	8		-56	PT condition holds good